OAR 340-245-8010
Table 1
Risk Action Levels†

Applicability	Risk Action Level	Excess Cancer Risk per Million	Noncancer Hazard Index
New and Reconstructed Source	Aggregate TEU Level	0.5	0.1
	Source Permit Level	0.5	0.5
	Community Engagement Level	5	1
	TLAER Level	10	1
	Permit Denial Level	25	1
Existing Source	Aggregate TEU Level	2.5	0.1
	Source Permit Level	5	0.5
	Community Engagement Level	25	1
	TBACT Level	50	5[a] or 3[b] or Risk Determination Ratio of 1.0[c]
	Risk Reduction Level	200	10[a] or 6[b] or Risk Determination Ratio of 2.0[c]
	Immediate Curtailment Level	500	20[a] or 12[b] or Risk Determination Ratio of 4.0[c]

Footnotes for OAR 340-245-8010 Table 1:

† Facility risk that is equal to or less than the values in the table is considered compliant with the Risk Action Level. Risk Action Levels are considered consistent with benchmarks in Oregon Laws 2018, chapter 102 (Senate Bill (SB) 1541 (2018)).

a) If all toxic air contaminants emitted by the source are identified as HI5 in OAR 340-247-8010, Table 2, and OAR 340-245-8010, Table 2.

b) If all toxic air contaminants emitted by the source are identified as HI3 in OAR 340-247-8010, Table 2, and OAR 340-245-8010, Table 2.

c) If toxic air contaminants emitted by the source include contaminants listed as both HI3 and HI5 in OAR 340-247-8010, Table 2, and OAR 340-245-8010, Table 2, and a Risk Determination Ratio is required to be calculated under OAR 340-245-0200.

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
75-07-0	Acetaldehyde		HI3	0.45	140	12	620	5.5	620	470
60-35-5	Acetamide			0.050		1.3		0.60		
67-64-1	Acetone		HI3		31,000		140,000		140,000	62,000
75-05-8	Acetonitrile		HI3		60		260		260	
107-02-8	Acrolein		HI5		0.35		1.5		1.5	6.9
79-06-1	Acrylamide	g	HI3	0.0059	6.0	0.062	26	0.12	26	
79-10-7	Acrylic acid		HI3		1.0		4.4		4.4	6,000
107-13-1	Acrylonitrile		HI3	0.015	5.0	0.38	22	0.18	22	220
309-00-2	Aldrin			0.00020		0.0053		0.0024		
107-05-1	Allyl chloride		HI3	0.17	1.0	4.3	4.4	2.0	4.4	
7429-90-5	Aluminum and compounds	l	HI5		5.0		22		22	
7664-41-7	Ammonia		HI3		500		2,200		2,200	1,200
62-53-3	Aniline		HI5	0.63	1.0	16	4.4	7.5	4.4	
7440-36-0	Antimony and compounds	l	HI3		0.30		1.3		1.3	1.0
140-57-8	Aramite			0.14		3.7		1.7		
7440-38-2	Arsenic and compounds	l	HI3	2.4E-05	0.00017	0.0013	0.0024	0.00062	0.0024	0.20
7784-42-1	Arsine		HI3		0.015		0.066		0.066	0.20
1332-21-4	Asbestos	I		4.3E-06		0.00011		5.2E-05		
103-33-3	Azobenzene			0.032		0.84		0.39		
71-43-2	Benzene		HI3	0.13	3.0	3.3	13	1.5	13	29
92-87-5	Benzidine (and its salts)	g		4.2E-06		4.4E-05		8.6E-05		
100-44-7	Benzyl chloride		HI3	0.020	1.0	0.53	4.4	0.24	4.4	240
7440-41-7	Beryllium and compounds	l	HI3	0.00042	0.0070	0.011	0.031	0.0050	0.031	0.020
111-44-4	*Bis*(2-chloroethyl) ether (BCEE)		HI3	0.0014		0.037		0.017		120
542-88-1	*Bis*(chloromethyl) ether		HI5	7.7E-05		0.0020		0.00092		1.4

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
117-81-7	*Bis*(2-ethylhexyl) phthalate (DEHP)	c		0.080		11		5.0		
75-25-2	Bromoform			0.91		24		11		
74-83-9	Bromomethane (methyl bromide)		HI3		5.0		22		22	3,900
106-94-5	1-Bromopropane (n-propyl bromide)		HI3	0.48	33	12	150	5.7	150	1,700
106-99-0	1,3-Butadiene		HI3	0.033	2.0	0.86	8.8	0.40	8.8	660
78-93-3	2-Butanone (methyl ethyl ketone)		HI3		5,000		22,000		22,000	5,000
78-92-2	*sec*-Butyl alcohol		HI3		30,000		130,000		130,000	
7440-43-9	Cadmium and compounds	c, l	HI3	0.00056	0.0050	0.014	0.037	0.0067	0.037	0.030
105-60-2	Caprolactam		HI3		2.2		9.7		9.7	50
75-15-0	Carbon disulfide		HI3		800		3,500		3,500	6,200
56-23-5	Carbon tetrachloride		HI3	0.17	100	4.3	440	2.0	440	1,900
463-58-1	Carbonyl sulfide		HI3		10		44		44	660
57-74-9	Chlordane		HI3	0.010	0.020	0.26	0.088	0.12	0.088	0.20
108171-26-2	Chlorinated paraffins	j		0.040		1.0		0.48		
7782-50-5	Chlorine		HI3		0.15		0.66		0.66	170
10049-04-4	Chlorine dioxide		HI3		0.60		2.6		2.6	2.8
532-27-4	2-Chloroacetophenone		HI5		0.030		0.13		0.13	
108-90-7	Chlorobenzene		HI3		50		220		220	
75-68-3	1-Chloro-1,1-difluoroethane		HI3		50,000		220,000		220,000	
75-45-6	Chlorodifluoromethane (Freon 22)		HI3		50,000		220,000		220,000	
75-00-3	Chloroethane (ethyl chloride)		HI3		30,000		130,000		130,000	40,000
67-66-3	Chloroform		HI3		300		1,300		1,300	490
74-87-3	Chloromethane (methyl chloride)		HI3		90		400		400	1,000

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
95-83-0	4-Chloro-*o*-phenylenediamine			0.22		5.7		2.6		
76-06-2	Chloropicrin		HI3		0.40		1.8		1.8	29
126-99-8	Chloroprene		HI3	0.0033	20	0.087	88	0.040	88	
95-69-2	*p*-Chloro-*o*-toluidine			0.013		0.34		0.16		
18540-29-9	Chromium VI, chromate and dichromate particulate	c, d	HI3	3.1E-05	0.083	0.00052	0.88	0.0010	0.88	0.30
7738-94-5	Chromium VI, chromic acid aerosol mist and chromium trioxide	c, d	HI3	3.1E-05	0.0021	0.00052	0.022	0.0010	0.022	0.0050
7440-48-4	Cobalt and compounds	l	HI3		0.10		0.44		0.44	
148[†]	Coke oven emissions	g		0.00095		0.0100		0.019		
7440-50-8	Copper and compounds	l	HI3							100
120-71-8	*p*-Cresidine			0.023		0.60		0.28		
1319-77-3	Cresols (mixture), including *m*-cresol, *o*-cresol, *p*-cresol		HI3		600		2,600		2,600	
135-20-6	Cupferron			0.016		0.41		0.19		
74-90-8	Cyanide, hydrogen		HI3		0.80		3.5		3.5	340
110-82-7	Cyclohexane		HI3		6,000		26,000		26,000	
50-29-3	DDT	e		0.010		0.27		0.12		
615-05-4	2,4-Diaminoanisole			0.15		3.9		1.8		
95-80-7	2,4-Diaminotoluene (2,4-toluene diamine)			0.00091		0.024		0.011		
333-41-5	Diazinon		HI3							10
96-12-8	1,2-Dibromo-3-chloropropane (DBCP)	g	HI3	9.8E-05	0.20	0.0010	0.88	0.0020	0.88	1.9
106-46-7	*p*-Dichlorobenzene (1,4-dichlorobenzene)		HI3	0.091	60	2.4	260	1.1	260	12,000
91-94-1	3,3'-Dichlorobenzidine			0.0029		0.076		0.035		
75-34-3	1,1-Dichloroethane (ethylidene dichloride)			0.63		16		7.5		

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACT RAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
156-60-5	*trans*-1,2-dichloroethene		HI3							790
75-09-2	Dichloromethane (methylene chloride)		HI3	59	600	620	2,600	1,200	2,600	2,100
78-87-5	1,2-Dichloropropane (propylene dichloride)		HI3		4.0		18		18	230
542-75-6	1,3-Dichloropropene		HI3	0.25	32	6.5	140	3.0	140	36
62-73-7	Dichlorvos (DDVP)		HI5		0.54		2.4		2.4	18
60-57-1	Dieldrin			0.00022		0.0057		0.0026		
200[†]	Diesel particulate matter		HI3	0.10	5.0	2.6	22	1.2	22	
111-42-2	Diethanolamine		HI3		0.20		0.88		0.88	
112-34-5	Diethylene glycol monobutyl ether		HI3		0.10		0.44		0.44	
111-90-0	Diethylene glycol monoethyl ether		HI5		0.30		1.3		1.3	
75-37-6	1,1-Difluoroethane		HI5		40,000		180,000		180,000	
60-11-7	4-Dimethylaminoazobenzene			0.00077		0.020		0.0092		
68-12-2	Dimethyl formamide		HI3		80		350		350	
57-14-7	1,1-Dimethylhydrazine		HI3							0.49
121-14-2	2,4-Dinitrotoluene			0.011		0.29		0.13		
123-91-1	1,4-Dioxane		HI3	0.20	30	5.2	130	2.4	130	7,200
122-66-7	1,2-Diphenylhydrazine (hydrazobenzene)			0.0045		0.12		0.055		
1937-37-7	Direct Black 38			7.1E-06		0.00019		8.6E-05		
2602-46-2	Direct Blue 6			7.1E-06		0.00019		8.6E-05		
16071-86-6	Direct Brown 95 (technical grade)			7.1E-06		0.00019		8.6E-05		
298-04-4	Disulfoton		HI3							6.0
106-89-8	Epichlorohydrin		HI3	0.043	3.0	1.1	13	0.52	13	1,300
106-88-7	1,2-Epoxybutane		HI5		20		88		88	
140-88-5	Ethyl acrylate		HI3		8.0		35		35	

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m^3)	(µg/m^3)	(µg/m^3)	(µg/m^3)	(µg/m^3)	(µg/m^3)	(µg/m^3)
100-41-4	Ethyl benzene		HI3	0.40	260	10	1,100	4.8	1,100	22,000
106-93-4	Ethylene dibromide (EDB, 1,2-dibromoethane)		HI3	0.0017	9.0	0.043	40	0.020	40	
107-06-2	Ethylene dichloride (EDC, 1,2-dichloroethane)		HI3	0.038	7.0	1.0	31	0.46	31	
107-21-1	Ethylene glycol		HI3		400		1,800		1,800	2,000
111-76-2	Ethylene glycol monobutyl ether		HI3		82		360		360	29,000
110-80-5	Ethylene glycol monoethyl ether		HI3		70		310		310	370
111-15-9	Ethylene glycol monoethyl ether acetate		HI3		60		260		260	140
109-86-4	Ethylene glycol monomethyl ether		HI3		60		260		260	93
110-49-6	Ethylene glycol monomethyl ether acetate		HI3		1.0		4.4		4.4	
75-21-8	Ethylene oxide	g	HI3	0.00020	30	0.0021	130	0.0040	130	160
96-45-7	Ethylene thiourea			0.077		2.0		0.92		
239[†]	Fluorides	c	HI3		2.3		20		20	240
7782-41-4	Fluorine gas		HI3							16
50-00-0	Formaldehyde		HI3	0.17	9.0	4.3	40	2.0	40	49
111-30-8	Glutaraldehyde		HI5		0.080		0.35		0.35	4.1
76-44-8	Heptachlor			0.00077		0.020		0.0092		
1024-57-3	Heptachlor epoxide			0.00038		0.010		0.0046		
118-74-1	Hexachlorobenzene			0.0020		0.051		0.024		
87-68-3	Hexachlorobutadiene			0.045		1.2		0.55		
608-73-1	Hexachlorocyclohexanes (mixture) including but not limited to:	c		0.00017		0.018		0.0084		
319-84-6	Hexachlorocyclohexane, *alpha*-	c		0.00017		0.018		0.0084		

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
319-85-7	Hexachlorocyclohexane, *beta*-	c		0.00017		0.018		0.0084		
58-89-9	Hexachlorocyclohexane, *gamma*- (Lindane)	c		0.00060		0.065		0.030		
77-47-4	Hexachlorocyclopentadiene		HI3		0.20		0.88		0.88	110
67-72-1	Hexachloroethane		HI3		30		130		130	58,000
822-06-0	Hexamethylene-1,6-diisocyanate		HI5		0.069		0.30		0.30	0.21
110-54-3	Hexane		HI3		700		3,100		3,100	
302-01-2	Hydrazine		HI3	0.00020	0.030	0.0053	0.13	0.0024	0.13	5.2
7647-01-0	Hydrochloric acid		HI3		20		88		88	2,100
7664-39-3	Hydrogen fluoride	c	HI3		2.1		19		19	16
7783-06-4	Hydrogen sulfide		HI3		2.0		8.8		8.8	98
78-59-1	Isophorone		HI3		2,000		8,800		8,800	
67-63-0	Isopropyl alcohol		HI3		200		880		880	3,200
98-82-8	Isopropylbenzene (cumene)		HI3		400		1,800		1,800	
7439-92-1	Lead and compounds	c, l	HI3		0.15		0.66		0.66	0.15
108-31-6	Maleic anhydride		HI5		0.70		3.1		3.1	
7439-96-5	Manganese and compounds	l	HI3		0.090		0.40		0.40	0.30
7439-97-6	Mercury and compounds	c, l	HI3		0.077		0.63		0.63	0.60
67-56-1	Methanol		HI3		4,000		18,000		18,000	28,000
101-14-4	4,4'-Methylene *bis*(2-chloroaniline) (MOCA)			0.0023		0.060		0.028		
101-77-9	4,4'-Methylenedianiline (and its dichloride)		HI5	0.00030	20	0.023	88	0.010	88	
101-68-8	Methylene diphenyl diisocyanate (MDI)		HI3		0.080		0.35		0.35	12
108-10-1	Methyl isobutyl ketone (MIBK, hexone)		HI3		3,000		13,000		13,000	
624-83-9	Methyl isocyanate		HI3		1.0		4.4		4.4	

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
80-62-6	Methyl methacrylate		HI5		700		3,100		3,100	
1634-04-4	Methyl *tert*-butyl ether		HI3	3.8	8,000	100	35,000	46	35,000	8,000
90-94-8	Michler's ketone			0.0040		0.10		0.048		
91-20-3	Naphthalene	c	HI3	0.029	3.7	0.76	16	0.35	16	200
365†	Nickel compounds, insoluble	f	HI3	0.0038	0.014	0.10	0.062	0.046	0.062	0.20
368†	Nickel compounds, soluble	f	HI3		0.014		0.062		0.062	0.20
7697-37-2	Nitric acid		HI5							86
98-95-3	Nitrobenzene		HI3	0.025	9.0	0.65	40	0.30	40	
79-46-9	2-Nitropropane		HI3		20		88		88	
924-16-3	N-Nitrosodibutylamine			0.00032		0.0084		0.0039		
55-18-5	N-Nitrosodiethylamine	g		5.9E-05		0.00062		0.0012		
62-75-9	N-Nitrosodimethylamine	g		0.00013		0.0013		0.0026		
86-30-6	N-Nitrosodiphenylamine			0.38		10		4.6		
156-10-5	*p*-Nitrosodiphenylamine			0.16		4.1		1.9		
621-64-7	N-Nitrosodipropylamine			0.00050		0.013		0.0060		
10595-95-6	N-Nitrosomethylethylamine			0.00016		0.0041		0.0019		
59-89-2	N-Nitrosomorpholine			0.00053		0.014		0.0063		
100-75-4	N-Nitrosopiperidine			0.00037		0.0096		0.0044		
930-55-2	N-Nitrosopyrrolidine			0.0017		0.043		0.020		
8014-95-7	Oleum (fuming sulfuric acid)		HI3							120
56-38-2	Parathion		HI3							0.020
87-86-5	Pentachlorophenol			0.20		5.1		2.4		
108-95-2	Phenol		HI3		200		880		880	5,800
75-44-5	Phosgene		HI3		0.30		1.3		1.3	4.0
7803-51-2	Phosphine		HI3		0.80		3.5		3.5	
7664-38-2	Phosphoric acid		HI5		10		44		44	
12185-10-3	Phosphorus, white		HI3		9.0		40		40	20

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
85-44-9	Phthalic anhydride		HI3		20		88		88	
447[†]	Polybrominated diphenyl ethers (PBDEs)	h	HI3							6.0
1336-36-3	Polychlorinated biphenyls (PCBs)	c		0.00053		0.020		0.0092		
645[†]	Polychlorinated biphenyls (PCBs) TEQ	c	HI3	1.0E-09	1.3E-07	9.0E-08	2.6E-05	4.2E-08	2.6E-05	
32598-13-3	PCB 77 [3,3',4,4'-tetrachlorobiphenyl]	c	HI3	1.0E-05	0.0013	0.00090	0.26	0.00042	0.26	
70362-50-4	PCB 81 [3,4,4',5-tetrachlorobiphenyl]	c	HI3	3.4E-06	0.00042	0.00030	0.085	0.00014	0.085	
32598-14-4	PCB 105 [2,3,3',4,4'-pentachlorobiphenyl]	c	HI3	3.4E-05	0.0042	0.0030	0.85	0.0014	0.85	
74472-37-0	PCB 114 [2,3,4,4',5-pentachlorobiphenyl]	c	HI3	3.4E-05	0.0042	0.0030	0.85	0.0014	0.85	
31508-00-6	PCB 118 [2,3',4,4',5-pentachlorobiphenyl]	c	HI3	3.4E-05	0.0042	0.0030	0.85	0.0014	0.85	
65510-44-3	PCB 123 [2,3',4,4',5'-pentachlorobiphenyl]	c	HI3	3.4E-05	0.0042	0.0030	0.85	0.0014	0.85	
57465-28-8	PCB 126 [3,3',4,4',5-pentachlorobiphenyl]	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	
38380-08-4	PCB 156 [2,3,3',4,4',5-hexachlorobiphenyl]	c	HI3	3.4E-05	0.0042	0.0030	0.85	0.0014	0.85	
69782-90-7	PCB 157 [2,3,3',4,4',5'-hexachlorobiphenyl]	c	HI3	3.4E-05	0.0042	0.0030	0.85	0.0014	0.85	
52663-72-6	PCB 167 [2,3',4,4',5,5'-hexachlorobiphenyl]	c	HI3	3.4E-05	0.0042	0.0030	0.85	0.0014	0.85	
32774-16-6	PCB 169 [3,3',4,4',5,5'-hexachlorobiphenyl]	c	HI3	3.4E-08	4.2E-06	3.0E-06	0.00085	1.4E-06	0.00085	
39635-31-9	PCB 189 [2,3,3',4,4',5,5'-heptachlorobiphenyl]	c	HI3	3.4E-05	0.0042	0.0030	0.85	0.0014	0.85	
646[†]	Polychlorinated dibenzo-*p*-dioxins (PCDDs) & dibenzofurans (PCDFs) TEQ	c	HI3	1.0E-09	1.3E-07	9.0E-08	2.6E-05	4.2E-08	2.6E-05	

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m^3)	(µg/m^3)	(µg/m^3)	(µg/m^3)	(µg/m^3)	(µg/m^3)	(µg/m^3)
1746-01-6	2,3,7,8-Tetrachlorodibenzo-*p*-dioxin (TCDD)	c	HI3	1.0E-09	1.3E-07	9.0E-08	2.6E-05	4.2E-08	2.6E-05	
40321-76-4	1,2,3,7,8-Pentachlorodibenzo-*p*-dioxin (PeCDD)	c	HI3	1.0E-09	1.3E-07	9.0E-08	2.6E-05	4.2E-08	2.6E-05	
39227-28-6	1,2,3,4,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD)	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	
57653-85-7	1,2,3,6,7,8-Hexachlorodibenzo-*p*-dioxin (HxCDD)	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	
19408-74-3	1,2,3,7,8,9-Hexachlorodibenzo-*p*-dioxin (HxCDD)	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	
35822-46-9	1,2,3,4,6,7,8-Heptachlorodibenzo-*p*-dioxin (HpCDD)	c	HI3	1.0E-07	1.3E-05	9.0E-06	0.0026	4.2E-06	0.0026	
3268-87-9	Octachlorodibenzo-*p*-dioxin (OCDD)	c	HI3	3.4E-06	0.00042	0.00030	0.085	0.00014	0.085	
51207-31-9	2,3,7,8-Tetrachlorodibenzofuran (TCDF)	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	
57117-41-6	1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	c	HI3	3.4E-08	4.2E-06	3.0E-06	0.00085	1.4E-06	0.00085	
57117-31-4	2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	c	HI3	3.4E-09	4.2E-07	3.0E-07	8.5E-05	1.4E-07	8.5E-05	
70648-26-9	1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	
57117-44-9	1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	
72918-21-9	1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
60851-34-5	2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	c	HI3	1.0E-08	1.3E-06	9.0E-07	0.00026	4.2E-07	0.00026	
67562-39-4	1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	c	HI3	1.0E-07	1.3E-05	9.0E-06	0.0026	4.2E-06	0.0026	
55673-89-7	1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	c	HI3	1.0E-07	1.3E-05	9.0E-06	0.0026	4.2E-06	0.0026	
39001-02-0	Octachlorodibenzofuran (OCDF)	c	HI3	3.4E-06	0.00042	0.00030	0.085	0.00014	0.085	
401†	Polycyclic aromatic hydrocarbons (PAHs)	c, g, n		4.3E-05		0.0016		0.0030		
191-26-4	Anthanthrene	c, g		0.00011		0.0039		0.0076		
56-55-3	Benz[a]anthracene	c, g		0.00021		0.0078		0.015		
50-32-8	Benzo[a]pyrene	c, g	HI3	4.3E-05	0.0020	0.0016	0.0088	0.0030	0.0088	0.0020
205-99-2	Benzo[b]fluoranthene	c, g		5.3E-05		0.0020		0.0038		
205-12-9	Benzo[c]fluorene	c, g		2.1E-06		7.8E-05		0.00015		
191-24-2	Benzo[g,h,i]perylene	c, g		0.0047		0.17		0.34		
205-82-3	Benzo[j]fluoranthene	c, g		0.00014		0.0052		0.010		
207-08-9	Benzo[k]fluoranthene	c, g		0.0014		0.052		0.10		
218-01-9	Chrysene	c, g		0.00043		0.016		0.030		
27208-37-3	Cyclopenta[c,d]pyrene	c, g		0.00011		0.0039		0.0076		
53-70-3	Dibenz[a,h]anthracene	c, g		4.3E-06		0.00016		0.00030		
192-65-4	Dibenzo[a,e]pyrene	c, g		0.00011		0.0039		0.0076		
189-64-0	Dibenzo[a,h]pyrene	c, g		4.7E-05		0.0017		0.0034		
189-55-9	Dibenzo[a,i]pyrene	c, g		7.1E-05		0.0026		0.0051		
191-30-0	Dibenzo[a,l]pyrene	c, g		1.4E-06		5.2E-05		0.00010		
206-44-0	Fluoranthene	c, g		0.00053		0.020		0.038		
193-39-5	Indeno[1,2,3-cd]pyrene	c, g		0.00061		0.022		0.043		

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
3697-24-3	5-Methylchrysene	c, g		4.3E-05		0.0016		0.0030		
7496-02-8	6-Nitrochrysene	c, g		4.3E-06		0.00016		0.00030		
7758-01-2	Potassium bromate			0.0071		0.19		0.086		
1120-71-4	1,3-Propane sultone			0.0014		0.038		0.017		
123-38-6	Propionaldehyde		HI5		8.0		35		35	
115-07-1	Propylene		HI5		3,000		13,000		13,000	
6423-43-4	Propylene glycol dinitrate		HI5		0.27		1.2		1.2	20
107-98-2	Propylene glycol monomethyl ether		HI3		7,000		31,000		31,000	
75-56-9	Propylene oxide		HI3	0.27	30	7.0	130	3.2	130	3,100
572[†]	Refractory ceramic fibers	i	HI5		0.030		0.13		0.13	
7783-07-5	Selenide, hydrogen		HI3							5.0
7782-49-2	Selenium and compounds	l	HI3							2.0
7631-86-9	Silica, crystalline (respirable)		HI5		3.0		13		13	
1310-73-2	Sodium hydroxide		HI3							8.0
100-42-5	Styrene		HI3		1,000		4,400		4,400	21,000
7664-93-9	Sulfuric acid		HI5		1.0		4.4		4.4	120
505-60-2	Sulfur mustard		HI3							0.70
7446-11-9	Sulfur trioxide		HI5		1.0		4.4		4.4	120
630-20-6	1,1,1,2-Tetrachloroethane			0.14		3.5		1.6		
79-34-5	1,1,2,2-Tetrachloroethane			0.017		0.45		0.21		
127-18-4	Tetrachloroethene (perchloroethylene)		HI3	3.8	41	100	180	46	180	41
811-97-2	1,1,1,2-Tetrafluoroethane		HI3		80,000		350,000		350,000	
62-55-5	Thioacetamide			0.00059		0.015		0.0071		
7550-45-0	Titanium tetrachloride		HI3		0.10		0.44		0.44	10
108-88-3	Toluene		HI3		5,000		22,000		22,000	7,500

OAR 340-245-8010
Table 2
Risk-Based Concentrations

			Non cancer TBACTRAL [m]	Residential Chronic		Non-Residential Chronic				Acute
				Cancer RBC[a]	Non-cancer RBC[a]	Child Cancer RBC[a]	Child Non-cancer RBC[a]	Worker Cancer RBC[a]	Worker Non-cancer RBC[a]	Non-cancer RBC[a]
CASRN[b]	Chemical	Notes		(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)
26471-62-5	Toluene diisocyanates (2,4- and 2,6-)		HI3	0.091	0.021	2.4	0.092	1.1	0.092	0.071
8001-35-2	Toxaphene (polychlorinated camphenes)			0.0031		0.081		0.038		
71-55-6	1,1,1-Trichloroethane (methyl chloroform)		HI3		5,000		22,000		22,000	11,000
79-00-5	1,1,2-Trichloroethane (vinyl trichloride)			0.063		1.6		0.75		
79-01-6	Trichloroethene (TCE, trichloroethylene)	g	HI3	0.20	2.1	3.5	9.2	2.9	9.2	2.1
88-06-2	2,4,6-Trichlorophenol			0.050		1.3		0.60		
96-18-4	1,2,3-Trichloropropane		HI5		0.30		1.3		1.3	1.8
121-44-8	Triethylamine		HI3		200		880		880	2,800
526-73-8	1,2,3-Trimethylbenzene		HI3		60		260		260	
95-63-6	1,2,4-Trimethylbenzene		HI3		60		260		260	
108-67-8	1,3,5-Trimethylbenzene		HI3		60		260		260	
51-79-6	Urethane (ethyl carbamate)	g		0.0020		0.021		0.041		
7440-62-2	Vanadium (fume or dust)		HI3		0.10		0.44		0.44	0.80
1314-62-1	Vanadium pentoxide		HI3	0.00012	0.0070	0.0031	0.031	0.0014	0.031	30
108-05-4	Vinyl acetate		HI3		200		880		880	200
593-60-2	Vinyl bromide		HI5		3.0		13		13	
75-01-4	Vinyl chloride	g, k	HI3	0.11	100	0.22	440	2.7	440	1,300
75-35-4	Vinylidene chloride		HI3		200		880		880	200
1330-20-7	Xylene (mixture), including *m*-xylene, *o*-xylene, *p*-xylene		HI3		220		970		970	8,700

Footnotes for OAR 340-245-8010 Table 2:

† Chemical designated by DEQ ID number.
a) RBC = Risk-Based Concentration
b) CASRN = Chemical Abstracts Service Registry Number, or DEQ ID if there is no CASRN.
c) Chronic RBCs include factors for multipathway risk.

d) The RBCs presented for chromium are applicable to hexavalent chromium. In the absence of data indicating otherwise, assume that any total chromium (i.e., unspeciated) that is measured or modeled is entirely in the hexavalent form. Determine, based on information about the source of emissions, whether hexavalent chromium is emitted in aerosol or particulate form, and apply the corresponding RBC. Because there are no RBCs for trivalent chromium, a source determined to be emitting only trivalent chromium cannot be shown to pose an unacceptable risk, so the risk in this case will be considered acceptable.

e) DDT RBCs apply to the sum of DDT, DDE, and DDD compounds.

f) As recommended by DEQ's Air Toxics Science Advisory Committee (ATSAC) in 2018, the two categories of nickel compounds contain the following specific nickel compounds:
Soluble nickel compounds are considered to be emitted mainly in aerosol form, to be less potent carcinogens than insoluble nickel compounds, and include nickel acetate, nickel chloride, nickel carbonate, nickel hydroxide, nickelocene, nickel sulfate, nickel sulfate hexahydrate, nickel nitrate hexahydrate, nickel carbonate hydroxide.
Insoluble nickel compounds are considered to be emitted mainly in particulate form, to be more potent carcinogens than soluble nickel compounds, and to include nickel subsulfide, nickel oxide, nickel sulfide, nickel metal.

g) RBCs adjusted to protect early-life exposure to infants and children because chemical is carcinogenic by a mutagenic mode of action.

h) RBCs apply to octabrominated diphenyl ethers (CASRN 32536-52-0) and pentabrominated diphenyl ethers (CASRN 32534-81-9), including BDE-99.

i) RBCs for asbestos and refractory ceramic fibers are in units of fibers/cm^3.

j) Chlorinated paraffins of average chain length of C12, approximately 60% chlorine by weight.

k) DEQ followed the ATSAC recommendation to develop a vinyl chloride TRV that already includes early-life exposure.

l) An inorganic chemical designated with "and compounds" indicates that the RBC applies to the sum of all forms of the chemical, expressed as the inorganic element.

m) Noncancer TBACT RAL = noncancer Toxics Best Available Control Technology Risk Action Level, OAR 340-245-8010, Table 1.

n) Because RBCs for PAHs were developed using TRVs for benzo[a]pyrene, apply PAH RBCs to summed benzo[a]pyrene toxicity equivalents for carcinogenic PAHs. If individual PAHs are not evaluated, apply PAH RBCs to total PAH concentrations.

OAR 340-245-8010 Table 3

Level 1 Risk Assessment Dispersion Factors

Table 3A: Stack Emission Dispersion Factors for Annual Exposure ($\mu g/m^3$ / pounds/year)

Stack	Exposure Location Distance (meters)												
Ht (m	50	60	70	80	90	100	110	120	130	140	150	160	170
5	0.0033	0.0026	0.0021	0.0017	0.0014	0.0012	0.0010	0.00088	0.00076	0.00066	0.00058	0.00051	0.00046
10	0.0014	0.0012	0.0011	0.00094	0.00084	0.00075	0.00068	0.00062	0.00057	0.00052	0.00048	0.00044	0.00041
15	0.00075	0.00061	0.00054	0.00049	0.00044	0.00040	0.00037	0.00034	0.00031	0.00029	0.00027	0.00025	0.00024
20	0.00072	0.00054	0.00035	0.00031	0.00028	0.00026	0.00023	0.00022	0.00020	0.00019	0.00017	0.00016	0.00015
25	0.00050	0.00041	0.00035	0.00025	0.00019	0.00018	0.00016	0.00015	0.00014	0.00013	0.00012	0.00012	0.00011
30	0.00037	0.00030	0.00026	0.00023	0.00019	0.00013	0.00012	0.00011	0.00010	0.000096	0.000090	0.000085	0.000080
35	0.00030	0.00023	0.00019	0.00017	0.00015	0.00013	0.00011	0.000081	0.000075	0.000071	0.000068	0.000064	0.000061
40	0.00023	0.00019	0.00015	0.00013	0.00012	0.00011	0.000096	0.000081	0.000064	0.000054	0.000051	0.000049	0.000047
45	0.00018	0.00016	0.00013	0.00011	0.000095	0.000085	0.000078	0.000072	0.000063	0.000053	0.000042	0.000038	0.000037
50	0.00014	0.00013	0.00011	0.000090	0.000077	0.000068	0.000062	0.000057	0.000053	0.000048	0.000042	0.000035	0.000029
Stack	Exposure Location Distance (meters)												
Ht (m	180	190	200	250	300	350	400	450	500	600	700	800	1000
5	0.00041	0.00037	0.00034	0.00023	0.00017	0.00013	0.00010	0.000084	0.000071	0.000052	0.000040	0.000032	0.000022
10	0.00038	0.00035	0.00033	0.00023	0.00017	0.00013	0.000098	0.000078	0.000064	0.000047	0.000036	0.000029	0.000021
15	0.00023	0.00021	0.00020	0.00016	0.00013	0.00010	0.000083	0.000069	0.000057	0.000041	0.000032	0.000025	0.000018
20	0.00014	0.00014	0.00013	0.00010	0.000086	0.000073	0.000062	0.000053	0.000046	0.000035	0.000027	0.000021	0.000015
25	0.00010	0.000096	0.000091	0.000072	0.000059	0.000051	0.000044	0.000039	0.000034	0.000027	0.000022	0.000018	0.000013
30	0.000075	0.000071	0.000068	0.000053	0.000044	0.000037	0.000032	0.000028	0.000025	0.000021	0.000017	0.000014	0.000010
35	0.000058	0.000055	0.000052	0.000042	0.000034	0.000029	0.000025	0.000022	0.000019	0.000016	0.000014	0.000011	0.000008
40	0.000045	0.000043	0.000041	0.000033	0.000028	0.000023	0.000020	0.000018	0.000016	0.000013	0.000011	0.000009	0.000007
45	0.000036	0.000034	0.000033	0.000027	0.000023	0.000019	0.000017	0.000015	0.000013	0.000011	0.000009	0.000008	0.000006
50	0.000027	0.000026	0.000026	0.000022	0.000019	0.000016	0.000014	0.000012	0.000011	0.000009	0.000007	0.000006	0.000005

Table 3B: Stack Emission Dispersion Factors for 24 hour Exposure ($\mu g/m^3$ / pounds/day)

Stack	Exposure Location Distance (meters)												
Ht (m	50	60	70	80	90	100	110	120	130	140	150	160	170
5	8.3	7.1	6.1	5.2	4.4	3.8	3.2	2.7	2.4	2.1	1.8	1.6	1.4
10	3.8	3.4	3.1	2.8	2.6	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5
15	1.8	1.6	1.6	1.5	1.4	1.3	1.2	1.1	1.1	1.00	0.95	0.91	0.87
20	1.6	1.3	0.91	0.86	0.82	0.77	0.73	0.69	0.65	0.62	0.59	0.56	0.54
25	0.97	0.93	0.85	0.64	0.52	0.50	0.48	0.46	0.44	0.42	0.40	0.38	0.36
30	0.62	0.59	0.57	0.55	0.49	0.34	0.32	0.31	0.30	0.29	0.28	0.27	0.26
35	0.42	0.41	0.39	0.38	0.37	0.34	0.29	0.22	0.21	0.21	0.20	0.20	0.19
40	0.30	0.29	0.28	0.28	0.27	0.26	0.25	0.22	0.17	0.15	0.15	0.15	0.14
45	0.22	0.22	0.21	0.21	0.20	0.20	0.19	0.19	0.17	0.16	0.12	0.11	0.11
50	0.16	0.16	0.16	0.16	0.16	0.15	0.15	0.14	0.14	0.13	0.12	0.10	0.082
Stack	Exposure Location Distance (meters)												
Ht (m	180	190	200	250	300	350	400	450	500	600	700	800	1000
5	1.3	1.2	1.1	0.72	0.55	0.44	0.36	0.30	0.26	0.20	0.16	0.13	0.092
10	1.4	1.3	1.3	0.91	0.67	0.50	0.38	0.30	0.25	0.18	0.14	0.12	0.088
15	0.83	0.80	0.77	0.64	0.53	0.43	0.36	0.30	0.25	0.18	0.13	0.10	0.075
20	0.52	0.49	0.48	0.40	0.35	0.31	0.27	0.23	0.20	0.16	0.12	0.096	0.064
25	0.35	0.34	0.32	0.27	0.23	0.21	0.19	0.17	0.15	0.12	0.100	0.082	0.057
30	0.25	0.24	0.23	0.19	0.17	0.15	0.13	0.12	0.11	0.095	0.078	0.066	0.048
35	0.18	0.18	0.17	0.15	0.13	0.11	0.099	0.090	0.083	0.072	0.062	0.053	0.040
40	0.14	0.14	0.13	0.11	0.10	0.088	0.078	0.070	0.064	0.056	0.049	0.044	0.033
45	0.11	0.11	0.10	0.092	0.081	0.072	0.065	0.058	0.053	0.045	0.040	0.036	0.028
50	0.081	0.080	0.079	0.072	0.065	0.059	0.053	0.048	0.044	0.037	0.032	0.029	0.024

Use of stack emission dispersion factors in a Level 1 screening risk assessment:

For each Toxics Emissions Unit, select the appropriate stack height and distance from the stack to nearest exposure locations approved by DEQ. For each exposure location, find the corresponding annual dispersion factor in Table 3A. For each toxic air contaminant, multiply the annual toxic air contaminant emission rate (in pounds/year) by the dispersion factor. Divide the product by the RBC for all the toxic air contaminants for the appropriate exposure location in OAR 340-245-8010 Table 2. Add up the resulting ratios for all Toxic Emissions Units for each exposure location. Compare the results with the Risk Action Levels in OAR 340-245-8010 Table 1. Repeat the process for daily emission rates (in pounds/day) using Table 3B at the acute exposure location.
For a stack height between the values shown in the table, either use the next lowest stack height, or interpolate the dispersion factor. For an exposure location distance between the values shown in the table, either use the next lowest distance, or interpolate the dispersion factor. For stack heights greater than 50 meters, use the appropriate dispersion factor for 50 meters. For exposure locations greater than 1,000 meters from the stack, use the appropriate dispersion factor at 1,000 meters. In the absence of a known stack height and exposure location distance, use as a default the annual dispersion factor (0.0033 $\mu g/m^3$ / pounds/year) and daily dispersion factor (8.3 $\mu g/m^3$ / pounds/day) for a stack height of 5 meters and an exposure location distance of 50 meters.
A Level 1 Risk Assessment will not be approved if the source is located near elevated terrain that DEQ determines could invalidate the assumptions used to develop the Level 1 Risk Assessment tool.

Stat. Auth.: ORS 468.020, 468.065, 468A.025, 468A.040, 468A.050, 468A.070, 468A.155
Stats. Implemented: ORS 468.065, 468A.010, 468A.015, 468A.025, 468A.035, 468A.040, 468A.050, 468A.070, and 468A.155

OAR 340-245-8010 Table 3

Level 1 Risk Assessment Dispersion Factors

Table 3C: Fugitive Emission Dispersion Factors for Annual Exposure ($\mu g/m^3$ / pounds/year)

Building Area	Building	Exposure Location Distance (meters)												
(1,000 ft2)	Height (ft)	50	60	70	80	90	100	110	120	130	140	150	160	170
≤3	≤20	0.0045	0.0033	0.0026	0.0020	0.0017	0.0014	0.0012	0.0010	0.00089	0.00078	0.00069	0.00062	0.00056
>3 to 6	≤20	0.0044	0.0032	0.0025	0.0020	0.0016	0.0014	0.0012	0.0010	0.00088	0.00077	0.00069	0.00061	0.00055
>3 to 6	>20	0.0041	0.0031	0.0024	0.0019	0.0016	0.0013	0.0011	0.0010	0.00086	0.00076	0.00067	0.00060	0.00054
>6 to 10	≤20	0.0044	0.0033	0.0025	0.0020	0.0017	0.0014	0.0012	0.0010	0.00088	0.00077	0.00069	0.00062	0.00055
>6 to 10	>20	0.0037	0.0028	0.0022	0.0018	0.0015	0.0013	0.0011	0.0010	0.00083	0.00074	0.00066	0.00059	0.00053
>10 to 15	≤20	0.0044	0.0033	0.0025	0.0020	0.0017	0.0014	0.0012	0.0010	0.00088	0.00077	0.00069	0.00062	0.00055
>10 to 15	>20	0.0034	0.0027	0.0021	0.0018	0.0015	0.0012	0.0011	0.00093	0.00081	0.00072	0.00064	0.00058	0.00052
>15 to 30	≤20	0.0043	0.0032	0.0025	0.0020	0.0016	0.0014	0.0012	0.0010	0.00088	0.00077	0.00069	0.00061	0.00055
>15 to 30	>20	0.0034	0.0027	0.0021	0.0018	0.0015	0.0012	0.0011	0.00093	0.00082	0.00072	0.00065	0.00058	0.00052
>30	>20	0.0022	0.0018	0.0015	0.0013	0.0011	0.0010	0.00086	0.00076	0.00068	0.00061	0.00055	0.00050	0.00046
Building Area	Building	Exposure Location Distance (meters)												
(1,000 ft2)	Height (ft)	180	190	200	250	300	350	400	450	500	600	700	800	1000
≤3	≤20	0.00050	0.00046	0.00042	0.00029	0.00021	0.00016	0.00013	0.00010	0.000087	0.000064	0.000049	0.000039	0.000027
>3 to 6	≤20	0.00050	0.00046	0.00042	0.00028	0.00021	0.00016	0.00013	0.00010	0.000087	0.000064	0.000049	0.000039	0.000027
>3 to 6	>20	0.00049	0.00045	0.00041	0.00028	0.00021	0.00016	0.00013	0.00010	0.000087	0.000064	0.000049	0.000039	0.000027
>6 to 10	≤20	0.00050	0.00046	0.00042	0.00028	0.00021	0.00016	0.00013	0.00010	0.000087	0.000064	0.000049	0.000039	0.000027
>6 to 10	>20	0.00048	0.00044	0.00041	0.00028	0.00020	0.00016	0.00013	0.00010	0.000086	0.000064	0.000049	0.000039	0.000027
>10 to 15	≤20	0.00050	0.00046	0.00042	0.00028	0.00021	0.00016	0.00013	0.00010	0.000087	0.000064	0.000049	0.000039	0.000027
>10 to 15	>20	0.00048	0.00044	0.00040	0.00028	0.00020	0.00016	0.00012	0.00010	0.000086	0.000063	0.000049	0.000039	0.000027
>15 to 30	≤20	0.00050	0.00046	0.00042	0.00028	0.00021	0.00016	0.00013	0.00010	0.000087	0.000064	0.000049	0.000039	0.000027
>15 to 30	>20	0.00048	0.00044	0.00040	0.00028	0.00020	0.00016	0.00013	0.00010	0.000086	0.000063	0.000049	0.000039	0.000027
>30	>20	0.00042	0.00039	0.00036	0.00025	0.00019	0.00015	0.00012	0.00010	0.000083	0.000061	0.000048	0.000038	0.000027

Table 3D: Fugitive Emission Dispersion Factors for 24 hour Exposure ($\mu g/m^3$ / pounds/day)

Building Area	Building	Exposure Location Distance (meters)												
(1,000 ft2)	Height (ft)	50	60	70	80	90	100	110	120	130	140	150	160	170
≤3	≤20	4.8	3.7	2.9	2.4	2.0	1.7	1.4	1.2	1.1	0.97	0.87	0.78	0.71
>3 to 6	≤20	4.1	3.1	2.5	2.0	1.7	1.4	1.2	1.1	0.95	0.84	0.76	0.68	0.62
>3 to 6	>20	3.5	2.8	2.2	1.9	1.6	1.3	1.2	1.0	0.90	0.80	0.72	0.65	0.59
>6 to 10	≤20	4.0	3.1	2.5	2.0	1.7	1.4	1.2	1.1	0.94	0.84	0.75	0.68	0.62
>6 to 10	>20	3.3	2.6	2.1	1.8	1.5	1.3	1.1	0.97	0.86	0.77	0.69	0.63	0.57
>10 to 15	≤20	4.0	3.1	2.4	2.0	1.7	1.4	1.2	1.1	0.94	0.84	0.75	0.68	0.62
>10 to 15	>20	2.9	2.4	2.0	1.6	1.4	1.2	1.1	0.93	0.83	0.74	0.67	0.61	0.56
>15 to 30	≤20	3.7	2.9	2.3	1.9	1.6	1.4	1.2	1.0	0.92	0.82	0.74	0.67	0.61
>15 to 30	>20	2.9	2.3	1.9	1.6	1.4	1.2	1.0	0.92	0.82	0.74	0.67	0.60	0.55
>30	>20	1.8	1.5	1.3	1.2	1.0	0.92	0.82	0.73	0.66	0.60	0.55	0.51	0.47
Building Area	Building	Exposure Location Distance (meters)												
(1,000 ft2)	Height (ft)	180	190	200	250	300	350	400	450	500	600	700	800	1000
≤3	≤20	0.65	0.59	0.55	0.38	0.29	0.22	0.18	0.15	0.13	0.095	0.074	0.060	0.043
>3 to 6	≤20	0.57	0.52	0.48	0.33	0.25	0.20	0.16	0.13	0.11	0.083	0.065	0.053	0.038
>3 to 6	>20	0.54	0.50	0.46	0.32	0.24	0.19	0.15	0.13	0.11	0.081	0.064	0.052	0.037
>6 to 10	≤20	0.56	0.52	0.48	0.33	0.25	0.20	0.16	0.13	0.11	0.083	0.065	0.053	0.038
>6 to 10	>20	0.53	0.48	0.45	0.31	0.24	0.19	0.15	0.12	0.11	0.080	0.063	0.051	0.036
>10 to 15	≤20	0.56	0.52	0.48	0.33	0.25	0.19	0.16	0.13	0.11	0.083	0.065	0.053	0.038
>10 to 15	>20	0.51	0.47	0.43	0.31	0.23	0.18	0.15	0.12	0.10	0.078	0.062	0.050	0.035
>15 to 30	≤20	0.55	0.51	0.47	0.33	0.25	0.19	0.16	0.13	0.11	0.083	0.065	0.053	0.037
>15 to 30	>20	0.51	0.47	0.43	0.31	0.23	0.18	0.15	0.12	0.10	0.078	0.062	0.050	0.035
>30	>20	0.43	0.40	0.37	0.27	0.21	0.17	0.14	0.12	0.098	0.075	0.059	0.048	0.034

Use of fugitive emission dispersion factors in a Level 1 screening risk assessment:

For each Toxics Emissions Unit, select the appropriate building dimensions and distance from building to nearest exposure locations approved by DEQ. For each exposure location, find the corresponding annual dispersion factor in Table 3C. For each toxic air contaminant, multiply the annual toxic air contaminant emission rate (in pounds/year) by the dispersion factor. Divide the product by the RBC for all the toxic air contaminants for the appropriate exposure location in OAR 340-245-8010 Table 2. Add up the resulting ratios for all Toxic Emissions Units for each exposure location. Compare the results with the Risk Action Levels in OAR 340-245-8010 Table 1. Repeat the process for daily emission rates (in pounds/day) using Table 3D at the acute exposure location.

For an exposure location distance between the values shown in the table, either use the next lowest distance, or interpolate the dispersion factor. For exposure locations greater than 1,000 meters from the building, use the appropriate dispersion factor at 1,000 meters. In the absence of known building dimensions and exposure location distance, use as a default, the annual dispersion factor (0.0045 $\mu g/m^3$ / pounds/year) and daily dispersion factor (4.8 $\mu g/m^3$ / pounds/day) for a building area of ≤3,000 ft2, height of ≤20 ft, and exposure location distance of 50 meters.

A Level 1 Risk Assessment will not be approved if the source is located near elevated terrain that DEQ determines could invalidate the assumptions used to develop the Level 1 Risk Assessment Dispersion Factors.

Stat. Auth.: ORS 468.020, 468.065, 468A.025, 468A.040, 468A.050, 468A.070, 468A.155
Stats. Implemented: ORS 468.065, 468A.010, 468A.015, 468A.025, 468A.035, 468A.040, 468A.050, 468A.070, and 468A.155